

RESEARCH ARTICLE

Open Access



The individualized selection of Pancreaticoenteric anastomosis in Pancreaticoduodenectomy

Ke-Min Jin[†], Wei Liu[†], Kun Wang, Quan Bao, Hong-Wei Wang and Bao-Cai Xing^{* ID}

Abstract

Background: The mortality following pancreaticoduodenectomy has markedly decreased but remains an important challenge for the complexity of operation and technical skills involved. The present study aimed to clarify the impact of individualized pancreaticoenteric anastomosis and management to postoperative pancreatic fistula.

Methods: Data from 529 consecutive pancreaticoduodenectomies were retrospectively analysed from the Hepatobiliary and Pancreatic Surgery Unit I, Peking Cancer Hospital. The pancreaticoenteric anastomosis was determined based on the pancreatic texture and diameter of the main pancreatic duct. The amylase value of the drainage fluid was dynamically monitored postoperatively on days 3, 5 and 7. A low speed intermittent irrigation was performed in selected patients. Intraoperative and postoperative results were collected and compared between the pancreaticogastrostomy (PG) group and pancreaticojejunostomy (PJ) group.

Results: From 2010 to 2019, 529 consecutive patients underwent pancreaticoduodenectomy. Pancreaticogastrostomy was performed in 364 patients; pancreaticojejunostomy was performed in 150 patients respectively. The clinically relevant pancreatic fistula (CR-POPF) was 9.8% and mortality was zero. The soft pancreas, diameter of main pancreatic duct ≤ 3 mm, BMI ≥ 25 , operation time > 330 min and pancreaticogastrostomy was correlated with postoperative pancreatic fistula significantly. The CR-POPF of PJ was significantly higher than that of PG in soft pancreas patients; the operation time of PJ was shorter than that of PG significantly in hard pancreas patients. Intraoperative blood loss and operation time of PG was less than that of PJ significantly in normal pancreatic duct patients ($p < 0.05$).

Conclusions: Individualized pancreaticoenteric anastomosis should be determined based on the pancreatic texture and pancreatic duct diameter. The appropriate anastomosis and postoperative management could prevent mortality.

Keywords: Pancreaticoduodenectomy anastomosis POPF

* Correspondence: xingbaocai88@sina.com

[†]Ke-Min Jin and Wei Liu contributed equally to this work.

Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Hepatobiliary and Pancreatic Surgery Unit I, Peking University Cancer Hospital & Institute, No. 52, Fu-Cheng Road, Beijing 100142, PR China



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Pancreaticoduodenectomy (PD) remains the golden standard for periampullary cancers [1]. Due to the complexity of the procedures and postoperative life-threatening complications, the mortality rates still was 1.4–29% [2, 3]. Despite a significant improvement in postoperative outcomes during recent decades, only a limited number of reports have documented zero mortality in consecutive pancreaticoduodenectomy series.

Postoperative pancreatic fistula (POPF) is one of the most potentially fatal complication after PD with rate ranging from 40 to 70%, which might cause arterial bleeding and mortality rate to 11–60% [4–6]. The established risk factors for POPF included a small pancreatic duct size, a soft pancreas and its posterior location. There have been a number of reported managements to reduce the incidence of POPF. The approach to management of the pancreatic remnant and form of pancreaticoenteric anastomosis (PA) determined the chance of developing POPF. Many efforts have been made to improve technical considerations through various modifications of pancreaticojejunostomy (PJ) and reconstruction with pancreaticogastrostomy (PG). The prompt management of POPF also decreased mortality, included in prophylactic use of octreotide and antibiotics.

Despite numerous trials comparing diverse PA techniques and other adjunctive strategies (pancreatic duct stenting, somatostatin analogues, etc), currently, there is no clear consensus regarding the ideal method of PA. The present study aimed to verify our management of individualized PA during operation and intermittent irrigation.

Methods

Study population

Between 1st November 2010 and 30th December 2019, 529 consecutive PDs were performed in the Hepatobiliary and Pancreatic Surgery Unit I of Peking Cancer Hospital. No 90-day mortality was reported in any of the 529 patients. Preoperative data were recorded, including demographics, comorbidity, performance status, American Society of Anaesthesiologists (ASA) level, previous abdominal history, preoperative serum bilirubin level and preoperative biliary drainage. The cases included 52 benign diseases (9.83%) and 477 malignancies (90.17%).

Surgical techniques

The texture of pancreatic parenchyma and the diameter of main pancreatic duct were assessed intraoperatively, which determined method of individualized PA. The texture was recorded as soft if elasticity of the pancreas was preserved (the stiffness of the patient's forehead as the reference as mentioned before). Additionally, the diameter of the main pancreatic duct was measured at

the surgically transected surface of the pancreas. PG was performed in patients with a normal main pancreatic duct (≤ 3 mm) and/or soft pancreas texture because the occurrence of clinically significant postoperative pancreatic fistula is more likely in such patients. It was performed with a two-layer purse-string suture on the posterior gastric wall, and the pancreas was mobilized 2–3 cm to be telescoped into the gastric cavity. PJ was performed in patients with the main duct was larger than 3 mm and hard pancreatic texture. It was performed with end-to-side two-layer sutures. The first layer was a duct-to-mucosa anastomosis, and the second-layer suture was located between the capsule of the pancreas and the seromuscular layer of the jejunum. A pancreatic duct stent was routinely used in all pancreaticodigestive reconstructions for internal drainage. For adequate drainage, two 20F Robinson drains were placed in addition to the pancreaticodigestive anastomosis.

Postoperative management

The amylase level of the drainage fluid was routinely evaluated on postoperative days 3, 5 and 7. POPF was defined as any measurable drainage on postoperative day 3 with an amylase content greater than 3 times the upper limit of the normal serum amylase level, and the severity was also graded as A, B, or C according to the criteria proposed by the International Study Group on Pancreatic Fistula⁶. If the fluid was transparent with normal amylase level, the drains were removed as soon as possible, typically on postoperative days 8. The drains were maintained for longer if patients had high drain amylase activity, copious fluid output or effluent with an unfavorable appearance (dark brown, greenish, milky or murky). Antibiotics were selected based on the susceptibility of bacteria isolated from drain fluids. At meanwhile, an irrigating tube was placed along the drainage tube and a low-speed intermittent irrigation was performed until the drain fluid returned transparent, which intended to dilute concentrated amylase. The speed of the irrigating saline should be controlled to avoid it spread into uninvolved areas of the abdominal cavity. When intra-abdominal collection and associated symptoms were detected, ultrasound-guided percutaneous drainage was performed. Irrigation was stopped, and the drainage was removed when the daily collections decreased < 30 mL/24 h. Postoperative complications were graded by the Clavien-Dindo classification [7].

Statistical analysis

Continuous variables were summarized as the medians and ranges, and categorical variables were summarized as frequencies and percentages. The Mann-Whitney U test was used to compare the continuous variables, and the Chi-square test was used to compare the categorical

Table 1 Demographic and comorbidity data

	Number(%)/median (range)
Age (years)	61(18–82)
Gender	
Male	294(55.6%)
Female	235(44.4%)
COPD	14(2.6%)
DM	94(17.8%)
HBP	151(28.5%)
PS	
0–1	501(94.7%)
2–3	28(5.3%)
ASA	
1	52(9.8%)
2	436(82.4%)
3	40(7.6%)
4	1(0.2%)
History of abdominal surgery	108(20.4%)
CAD	26(4.9%)
Cerebral vessel disease	23(4.3%)
Previous chemotherapy or radiotherapy	20(3.8%)
Preoperative biliary drainage	157(29.7%)
Preoperative hyperbilirubinemia	246(46.5%)
Weight loss	290(54.8%)
Smoking	162(30.6%)
Alcohol abuse	105(19.8%)

COPD chronic obstructive pulmonary disease, HBP hypertension, DM diabetes mellitus, PS performance status, CAD coronary artery disease, ASA American Society of Anaesthesiologists

variables. A *p* value of < 0.05 was deemed statistically significant.

Results

Patient characteristics and surgical details

The demographic and comorbidity data were shown in Table 1. The median age of all patients was 61 years old (range: 18–82). Two hundred and ninety-four patients (55.6%) were male. The most common disease was pancreatic ductal adenocarcinoma (130 cases, 24.6%) (Supplementary Table 1). The median operation time was 287 min (range: 150–648 min), and the median intraoperative blood loss was 200 ml (range: 50–1500 ml). The median postoperative length of the hospital stay was 19 days (range: 8–121 days) (Table 2).

Postoperative complications

As shown in Table 2, postoperative complications according to the Clavien-Dindo classification system

Table 2 Intraoperative variables and postoperative complications

Variables	Median (range)/number(%)
Operation time (min)	287(150–648)
Intraoperative blood loss (ml)	200(50–1500)
Transfusion	68(12.9%)
RBC transfused(U)	4(2–20)
Pancreatic remnant anastomosis	
PJ	150(28.4%)
PG	364(68.8%)
No anastomosis	15(2.8%)
Combined PV-SMV or IVC resection	20(3.8%)
Operation types	
PD	436(82.4%)
PD combined devisceration	78(14.7%)
TP	10(1.9%)
TP combined devisceration	5(0.9%)
Morbidity	311(58.8%)
Clavien-Dindo classification	
I	51(9.6%)
II	183(34.6%)
IIIa	56(10.6%)
IIIb	11(2.1%)
IVa	10(1.9%)
DGE	77(14.6%)
B	22(4.2%)
C	55(10.4%)
PPH	77(14.6%)
A	4(0.8%)
B	63(11.9%)
C	10(1.9%)
Bile leakage	21(4.0%)
Pneumonia	12(2.3%)
Abdominal infection	84(15.9%)
Urinary tract infection	3(0.6%)
Wound infection	5(0.9%)
Arrhythmia	10(1.9%)
Postoperative pancreatic fistula	204(38.6%)
A	152(28.7%)
B	46(8.7%)
C	6(1.1%)
Percutaneous drainage	53(10.0%)
RBC transfusion	124(23.4%)
RBC transfused(U)	4(2–30)
Relaparotomy	16(3.0%)
Readmission	6(1.1%)
Postoperative hospital stay (days)	19(8–121)

PV-SMV portal vein-superior mesenteric vein, IVC inferior vena cava, PD pancreaticoduodenectomy, TP total pancreatectomy, DGE delayed gastric emptying, PPH postpancreatectomy haemorrhage, RBC red blood cell

occurred in 311 patients (58.8%). POPF occurred in 204 patients (38.6%), with grade A POPF in 152 patients (28.7%), grade B POPF in 46 patients (8.7%), and grade C POPF in 6 patients (1.1%). POPF relating abdominal hemorrhage developed in 9 patients. Among of them, 3 patients underwent angiography and embolization, 5 patients underwent reoperation and one patient received conservative treatment. Twenty-one patients (4.0%) experienced bile leakage after the operation. Abdominal infection occurred in 84 patients (15.9%) and 53 patients (10.0%) received percutaneous drainage. One hundred and twenty-four patients (23.4%) received a postoperative RBC transfusion, and the median number of transfused RBCs was 4 units (range: 2–30 units).

PG versus PJ

The treatment groups were balanced in terms of most clinical factors and surgical details. However, the hard pancreas (PJ vs. PG, 88.0% vs. 15.4%, $p < 0.001$) and dilated pancreatic duct rates (PJ vs. PG, 65.3% vs. 13.5%, $p < 0.001$) were significantly different between the 2 groups (Table 3). The PJ group patients also had a significantly higher percentage of pancreatic ductal adenocarcinoma or chronic pancreatitis than the PG group (50.0% vs. 18.1%, $p < 0.001$). The incidence of POPF was 42.6% after PG and 32.7% after PJ ($p = 0.037$). Soft pancreas texture, a normal pancreatic duct size, obesity, longer operation time and pancreaticogastrostomy were significant factors affecting POPF, with estimated odds ratios of 3.191 ($p < 0.001$), 3.928 ($p < 0.001$), 1.976 ($p = 0.001$), 1.635 ($p = 0.040$) and 0.323 ($p = 0.001$) respectively (Table 4). In the subgroup analysis according to pancreas texture, the CR-POPF of PJ was higher than that of PG significantly in patients with soft pancreas texture ($p = 0.033$). The operation time of PJ was shorter than that of PG significantly in patients with hard pancreas texture ($p = 0.035$) (Table 5). In the subgroup analysis according to pancreatic duct size, intraoperative blood loss and operation time of PG was less than that of PJ significantly in patients with normal main pancreatic duct ($p < 0.05$) (Table 6).

Discussion

The present study reported 529 consecutive PDs, representing the largest consecutive study of PD without mortality. Although the occurrence of pancreatic fistulas was 38.6%, routine evaluation of amylase level of the drainage fluid and intermittent irrigation through drainage tube might prevent POPF related mortality. Recent advances in surgical techniques and adequate management of postoperative complications have led to improved clinical outcomes of PD, and the mortality following PD has decreased to below 6% [8]. POPF is a main source of major morbidity due to the intraperitoneal release of

Table 3 Comparison of the perioperative variables of the PJ and PG groups

	PJ(150)	PG(364)	p
MPD diameter(> 3 mm)	98(65.3%)	49(13.5%)	< 0.001
Pancreatic texture (hard)	132(88.0%)	56(15.4%)	< 0.001
Gender (male)	89(59.3%)	196(53.8%)	0.255
Pathology (PDAC or CP)	75(50.0%)	66(18.1%)	< 0.001
BMI	23.5 ± 3.2	23.6 ± 3.3	0.628
ASA			0.329
1	10(6.7%)	40(11.0%)	
2	131(87.3%)	294(80.8%)	
3	8(6.0%)	29(8.0%)	
4	0	1(0.3%)	
COPD	3(2.0%)	11(3.0%)	0.767
DM	27(18.0%)	61(16.8%)	0.734
HBP	45(30.0%)	101(27.7%)	0.607
History of abdominal surgery	35(23.3%)	67(18.4%)	0.203
Preoperative biliary drainage	50(33.3%)	104(28.6%)	0.284
Preoperative hyperbilirubinemia	87(58.0%)	157(43.1%)	0.002
Operation time (min)	296.4 ± 58.9	288.7 ± 65.3	0.211
Intraoperative blood loss (ml)	276.0 ± 186.2	236.4 ± 169.8	0.010
Postoperative hospital stay (days)	21.6 ± 12.2	24.2 ± 14.4	0.056
Complications	78(52.0%)	229(62.9%)	0.022
POPF	49(32.7%)	155(42.6%)	0.037
Grade A POPF	32(21.3%)	120(33.0%)	0.009
Grade B/C POPF	17(11.3%)	35(9.6%)	0.557
Relaparotomy	1(0.7%)	15(4.1%)	0.048
DGE	19(12.7%)	58(15.9%)	0.345
PPH	23(15.3%)	51(14.0%)	0.698

PJ pancreaticojejunostomy, *PG* pancreaticogastrostomy, *MPD* main pancreatic duct, *PDAC* pancreatic ductal adenocarcinoma, *BMI* body mass index, *ASA* American Society of Anaesthesiologists, *COPD* chronic obstructive pulmonary disease, *DM* diabetes mellitus, *HBP* hypertension, *DGE* delayed gastric emptying, *PPH* postpancreatectomy haemorrhage, *CP* chronic pancreatitis, *POPF* Postoperative pancreatic fistula

enterokinase and activation of pancreatic proenzymes resulting in sepsis and haemorrhage. This complication might be inevitable and still causes troublesome short-term outcomes after surgery.

The approach to management PA remains key factor in determining the chance of developing a POPF. Despite multiple randomized studies and meta-analyses, there is no clear evidence or universally accepted guidelines for how to construct the optimal PA after PD. [9] The multiple studies described above have failed to provide definitive, consistent, and convincing level 1 evidence that any one technique of PA is better than the others, either during the traditional open PD or more recently with the laparoscopic PD. Therefore, it should be

Table 4 Logistic Regression Analysis of factors affecting POPF

Parameter	Univariate Analysis		P	Multivariate Analysis		P
	Odds Ratio	95% CI		Odds Ratio	95% CI	
Gender			0.153			
Female	1					
Male	0.771	0.539–1.102				
Gland texture			< 0.001			< 0.001
Hard	1			1		
Soft	2.787	1.877–4.138		3.191	1.695–6.008	
Pathology			< 0.001			0.15
PDAC or CP	1			1		
Others	2.635	1.704–4.077		1.605	0.842–3.058	
pancreatic duct diameter			< 0.001			< 0.001
> 3 mm	1			1		
≤ 3 mm	4.14	2.601–6.591		3.928	2.217–6.961	
Intraoperative blood loss			0.814			
≤ 400 ml	1					
> 400 ml	1.076	0.582–1.992				
Age			0.725			
< 70	1					
≥ 70	0.913	0.550–1.516				
BMI			< 0.001			0.001
< 25	1			1		
≥ 25	2.068	1.421–3.008		1.976	1.326–2.945	
ASA score			0.4			
1-2	1					
3-4	0.743	0.303–1.134				
Preoperative jaundice			0.608			
No	1					
Yes	0.912	0.640–1.299				
Alcohol abuse			0.482			
No	1					
Yes	1.17	0.755–1.813				
Diabetes mellitus			0.462			
No	1					
Yes	1.19	0.748–1.894				
Previous abdominal surgery history			0.913			
No	1					
Yes	0.976	0.626–1.521				
Preoperative biliary decompression			0.176			
No	1					
Yes	1.302	0.888–1.910				
Operation time			0.028			0.04
≤ 330 min	1			1		
> 330 ml	1.622	1.055–2.494		1.635	1.022–2.617	
Intraoperative RBC transfusion			0.273			

Table 4 Logistic Regression Analysis of factors affecting POPF (Continued)

Parameter	Univariate Analysis		P	Multivariate Analysis		P
	Odds Ratio	95% CI		Odds Ratio	95% CI	
No	1					
Yes	1.345	0.791–2.286				
Pancreicoenteric anastomosis method			0.037			0.001
Pancreicojejunostomy (PJ)	1			1		
Pancrecicogastrostomy (PG)	1.529	1.025–2.279		0.323	0.163–0.639	
Combined with vascular resection			0.232			
No	1					
Yes	0.531	0.188–1.498				

PDAC pancreatic ductal adenocarcinoma;CP: chronic pancreatitis, BMI body weight index, ASA American Society of Anaesthesiologists

expected to utilize different forms of PA depending on pancreatic texture and main pancreatic duct in selected situations, which might be a potential solution to evade problem of POPF. PJ is the commonly preferred anastomosis method. Many techniques have been proposed for the reconstruction of pancreatic digestive continuity to prevent complications after PD. [10–12] PG anastomosis has an excellent blood supply, less tension in the anastomosis, and a thick stomach wall, which facilitate the establishment of a sound anastomosis [13]. Furthermore, the acid milieu of the stomach and the absence of enterokinase protect the anastomosis from autodigestion by inactivating the pancreatic proenzymes [14]. Previous studies reported contradictory results regarding the impact of PG versus PJ on the postoperative fistula rate [15–17]. Recently, reconstruction by PG was associated

with lower postoperative pancreatic and biliary fistula rates [18]. These principles include good exposure and visualization, the use of a fine, nonstrangulating suture to produce a water-tight patent anastomosis, preservation of the blood supply, tension-free fixation of the gastrointestinal tract to the pancreas, and coverage of the transected pancreas [19]. The present study identified the morbidity of GJ was higher than that of PG significantly for patients with soft pancreas. Generally, it recommended that PG was an optimal approach for these patients.

This study contained 52 patients (9.8%) with clinically relevant PF after PD, which was highly consistent with previous studies. A grade A POPF was not considered clinically important; thus, only grade B/C should account for the incidence of clinically relevant POPF [20].

Table 5 Subgroup analysis according to pancreas texture: PG vs PJ

	For hard pancreas texture			For soft pancreas texture		
	PJ(132pts)	PG(56pts)	p	PJ(18pts)	PG(308pts)	p
Operation time (min)	293.0 ± 56.1	312.6 ± 62.0	0.035	321.3 ± 74.1	284.4 ± 65.0	0.021
Intraoperative blood loss (ml)	270.8 ± 191.7	253.6 ± 151.9	0.550	313.9 ± 137.0	233.3 ± 172.9	0.053
POPF	28.8%	16.1%	0.066	61.1%	47.4%	0.258
CR-POPF	9.1%	1.8%	0.071	27.8%	11.0%	0.033
Intraoperative RBC transfusion	12.9%	12.5%	0.943	16.7%	11.7%	0.527
PPH	14.4%	16.1%	0.768	22.2%	13.6%	0.298
Postoperative RBC transfusion	19.7%	21.4%	0.787	50.0%	23.7%	0.012
Postoperative hospital stay (days)	20.1 ± 11.3	22.4 ± 10.2	0.207	32.5 ± 12.8	24.5 ± 15.0	0.028
Postoperative complications	48.5%	53.6%	0.524	77.8%	64.6%	0.315
Reoperation	0.0%	1.8%	0.298	5.6%	4.5%	0.582
Postoperative percutaneous drainage	6.1%	5.4%	0.851	16.7%	12.7%	0.714
Dilated main pancreatic duct(>3 mm)	71.2%	39.3%	< 0.001	22.2%	8.8%	0.079
Readmission	1.5%	0.0%	1.000	0.0%	1.3%	1.000
Postoperative abdominal infection	9.1%	10.7%	0.729	27.8%	19.8%	0.413

PG pancreaticogastrostomy, PJ pancreicojejunostomy, POPF postoperative pancreatic fistula, CR-POPF clinically relevant postoperative pancreatic fistula, PPH post-pancreatectomy hemorrhage, RBC red blood cell

Table 6 Subgroup analysis according to main pancreatic duct diameter: PG vs PJ

	For dilated main pancreatic duct(> 3 mm)			For normal main pancreatic duct(≤3 mm)		
	PJ(98pts)	PG(49pts)	<i>p</i>	PJ(52pts)	PG(315pts)	<i>p</i>
Operation time (min)	290.3 ± 58.3	300.5 ± 61.6	0.328	307.9 ± 59.0	286.9 ± 65.7	0.031
Intraoperative blood loss (ml)	264.3 ± 193.7	277.6 ± 160.1	0.680	298.1 ± 170.6	230.0 ± 170.6	0.008
POPF	20.4%	14.3%	0.366	55.8%	47.0%	0.240
CR-POPF	7.1%	2.0%	0.269	19.2%	10.8%	0.083
Intraoperative RBC transfusion	11.2%	12.2%	0.855	17.3%	11.7%	0.262
PPH	14.3%	10.2%	0.487	17.3%	14.6%	0.613
Postoperative RBC transfusion	17.3%	20.4%	0.651	34.6%	23.8%	0.097
Postoperative hospital stay (days)	18.4 ± 10.3	18.4 ± 6.9	0.985	27.8 ± 13.0	25.1 ± 15.1	0.228
Postoperative complications	39.8%	44.9%	0.554	75.0%	65.7%	0.187
Reoperation	0.0%	0.0%	–	1.9%	4.8%	0.711
Postoperative percutaneous drainage	5.1%	4.1%	1.000	11.5%	12.7%	0.815
Hard pancreatic gland texture	95.9%	44.9%	< 0.001	73.1%	10.8%	< 0.001
Readmission	1.5%	0.0%	1.000	0.0%	1.0%	1.000
Postoperative abdominal infection	2.0%	2.0%	1.000	21.2%	20.3%	0.890

PG pancreaticogastrostomy, PJ pancreaticojejunostomy, POPF postoperative pancreatic fistula, CR-POPF clinically relevant postoperative pancreatic fistula, PPH post-pancreatectomy hemorrhage, RBC red blood cell

The prevention of clinically relevant POPF may partially depend on the prompt healing of pancreaticodigestive tract anastomoses, which is attributed to the immediate recovery from a minor pancreatic fistula originating from the pancreatic branch duct or parenchyma at the pancreas surface [5]. Prophylactic drains after pancreatic surgery allow physicians to monitor the occurrence of intra-abdominal bleeding and to detect and drain a pancreatic, biliary, or enteric fistula [21]. Intermittent irrigation aimed to dilute the concentration of intra-abdominal amylase level, which is effective in preventing damage caused by the erosive retention of pancreatic secretions. For symptomatic abdominal collection fluid or abdominal abscess, percutaneous puncture and drainage was the preferred procedure. The reported success rate of the conservative treatment of a POPF is approximately 80%. The relaparotomy should be performed only when patients presented a high output fistula and severe sepsis or haemorrhage and cannot be managed by other means [22].

The results were consistent with the literature that reported a significantly elevated risk of post-PD bleeding in patients with pancreatic fistulas [23, 24]. It was also confirmed that pancreatic leakage and intra-abdominal abscess were correlated to post-PD bleeding [25]. Therefore, any procedure that can prevent pancreatic fistula or intra-abdominal abscess can decrease the post-PD bleeding rate. Prophylactic irrigation around a PJ was reported to possibly decrease the incidence of pancreatic fistulas and infectious complications [26]. It was routinely performed a low-

speed intermittent irrigation was added when the drain fluid turned turbid with sediment.

Conclusions

Individualized pancreatocentric anastomosis should be determined based on the pancreatic texture and diameter of the main pancreatic duct. The appropriate anastomosis and postoperative management could prevent mortality.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12893-020-00791-y>.

Additional file 1.

Abbreviations

CR-POPF: Clinically relevant postoperative pancreatic fistula; PG: Pancreaticogastrostomy; PJ: Pancreaticojejunostomy; BMI: Body mass index; PD: Pancreaticoduodenectomy; POPF: Postoperative pancreatic fistula; PA: Pancreatocentric anastomosis; ASA: American Society of Anaesthesiologists

Acknowledgements

Not applicable.

Authors' contributions

BCX contributed to the conception and design. KW, QB and HWW are responsible for the provision of the study materials and data collection. KMJ and WL contributed to the data analysis and interpretation and draft writing equally. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to protecting individual patient privacy but are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This study was approved by the Clinical Research Ethics Committee of the Peking University Cancer Hospital and was performed in compliance with the Helsinki Declaration. Written informed consent was obtained from all patients.

Consent for publication

Not Applicable.

Competing interests

The authors declare that they have no conflict of interest.

Received: 19 May 2020 Accepted: 9 June 2020

Published online: 22 June 2020

References

- He J, Ahuja N, Makary MA, et al. 2564 resected periampullary adenocarcinomas at a single institution: trends over three decades. *HPB (Oxford)*. 2014;16:83–90.
- Cameron JL, He J. Two thousand consecutive pancreaticoduodenectomies. *J Am Coll Surg*. 2015;220:530–6.
- Kawakatsu S, Inoue Y, Mise Y, et al. Comparison of pancreatojejunostomy techniques in patients with a soft pancreas: Kakita anastomosis and Blumgart anastomosis. *BMC Surg*. 2018;18:88.
- Hanaki T, Uejima C, Amisaki M, et al. The attenuation value of preoperative computed tomography as a novel predictor for pancreatic fistula after pancreaticoduodenectomy. *Surg Today*. 2018;48:598–608.
- Takahashi S, Gotohda N, Kato Y, et al. Measure of pancreas transection and postoperative pancreatic fistula. *J Surg Res*. 2016;202:276–83.
- Miura F, Asano T, Amano H, et al. Management of postoperative arterial hemorrhage after pancreato-biliary surgery according to the site of bleeding: re-laparotomy or interventional radiology. *J Hepato-Biliary-Pancreat Surg*. 2009;16:56–63.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240:205–13.
- Blanc T, Cortes A, Goere D, et al. Hemorrhage after pancreaticoduodenectomy: when is surgery still indicated? *Am J Surg*. 2007;194:3–9.
- Shrikhande SV, Sivasanker M, Vollmer CM, et al. Pancreatic anastomosis after pancreatoduodenectomy: a position statement by the international study Group of Pancreatic Surgery (ISGPS). *Surgery*. 2017;161:1221–34.
- Sakorafas GH, Friess H, Balsiger BM, et al. Problems of reconstruction during pancreatoduodenectomy. *Dig Surg*. 2001;18:363–9.
- Yeo CJ, Cameron JL, Lillemoe KD, et al. Does prophylactic octreotide decrease the rates of pancreatic fistula and other complications after pancreaticoduodenectomy? Results of a prospective randomized placebo-controlled trial. *Ann Surg*. 2000;232:419–29.
- Peng SY, Wang JW, Lau WY, et al. Conventional versus binding pancreatojejunostomy after pancreaticoduodenectomy: a prospective randomized trial. *Ann Surg*. 2007;245:692–8.
- Aranha GV, Hodul P, Golts E, et al. A comparison of pancreaticogastrostomy and pancreatojejunostomy following pancreaticoduodenectomy. *J Gastrointest Surg*. 2003;7:672–82.
- Fang WL, Shyr YM, Su CH, et al. Comparison between pancreatojejunostomy and pancreaticogastrostomy after pancreaticoduodenectomy. *J Formos Med Assoc*. 2007;106:717–27.
- Wellner U, Makowiec F, Fischer E, et al. Reduced postoperative pancreatic fistula rate after pancreatogastrostomy versus pancreatojejunostomy. *J Gastrointest Surg*. 2009;13:745–51.
- Wente MN, Shrikhande SV, Muller MW, et al. Pancreatojejunostomy versus pancreaticogastrostomy: systematic review and meta-analysis. *Am J Surg*. 2007;193:171–83.
- Fathy O, Abdel-Wahab M, Elghwalby N, et al. Surgical management of periampullary tumors: a retrospective study. *Hepatogastroenterology*. 2008;55:1463–9.
- Menahem B, Guittet L, Mulliri A, et al. Pancreaticogastrostomy is superior to pancreatojejunostomy for prevention of pancreatic fistula after pancreaticoduodenectomy: an updated meta-analysis of randomized controlled trials. *Ann Surg*. 2005;61:882–7.
- Adams DB. The pancreatic anastomosis: the danger of a leak, which anastomotic technique is better? *J Gastrointest Surg*. 2009;13:1182–3.
- Pedrazzoli S, Liessi G, Pasquali C, et al. Postoperative pancreatic fistulas: preventing severe complications and reducing reoperation and mortality rate. *Ann Surg*. 2009;249:97–104.
- Oguro S, Yoshimoto J, Imamura H, et al. Three hundred and sixty-eight consecutive pancreaticoduodenectomies with zero mortality. *J Hepatobiliary Pancreat Sci*. 2017;24:226–34.
- Alexakis N, Sutton R, Neoptolemos JP. Surgical treatment of pancreatic fistula. *Dig Surg*. 2004;21:262–74.
- Yekebas EF, Wolfram L, Cataldegirmen G, et al. Postpancreatectomy hemorrhage: diagnosis and treatment: an analysis in 1669 consecutive pancreatic resections. *Ann Surg*. 2007;246:269–80.
- Choi SH, Moon HJ, Heo JS, et al. Delayed hemorrhage after pancreaticoduodenectomy. *J Am Coll Surg*. 2004;199:186–91.
- Wei HK, Wang SE, Shyr YM, et al. Risk factors for post-pancreaticoduodenectomy bleeding and finding an innovative approach to treatment. *Dig Surg*. 2009;26:297–305.
- Nakano H, Asakura T, Sakurai J, et al. Prophylactic irrigation around a pancreatojejunostomy for the treatment of a pancreatic fistula after a pancreaticoduodenectomy in patients with a risky pancreatic remnant. *Hepatogastroenterology*. 2008;55:717–21.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

