


RESEARCH ARTICLE

Open Access



# Assessment of pain and postoperative nausea and vomiting and their association in the early postoperative period: an observational study from Palestine

Reem M. Elsaid<sup>1</sup>, Ashraquat S. Namrouti<sup>1</sup>, Ahmad M. Samara<sup>1</sup>, Wael Sadaqa<sup>1,2</sup> and Sa'ed H. Zyoud<sup>3,4,5\*</sup> 

## Abstract

**Background:** Postoperative nausea and vomiting (PONV) and postoperative pain (POP) are most commonly experienced in the early hours after surgery. Many studies have reported high rates of PONV and POP, and have identified factors that could predict the development of these complications. This study aimed to evaluate the relationship between PONV and POP, and to identify some factors associated with these symptoms.

**Methods:** This was a prospective, multicentre, observational study performed at An-Najah National University Hospital and Rafidia Governmental Hospital, the major surgical hospitals in northern Palestine, from October 2019 to February 2020. A data collection form, adapted from multiple previous studies, was used to evaluate factors associated with PONV and POP in patients undergoing elective surgery. Patients were interviewed during the first 24 h following surgery. Multiple binary logistic regression was applied to determine factors that were significantly associated with the occurrence of PONV.

**Results:** Of the 211 patients included, nausea occurred in 43.1%, vomiting in 17.5%, and PONV in 45.5%. Multiple binary logistic regression analysis, using PONV as a dependent variable, showed that only patients with a history of PONV [odds ratio (OR) = 2.28; 95% confidence interval (CI) = 1.03–5.01;  $p = 0.041$ ] and POP (OR = 2.41; 95% CI = 1.17–4.97;  $p = 0.018$ ) were significantly associated with the occurrence of PONV. Most participants (74.4%) reported experiencing pain at some point during the first 24 h following surgery. Additionally, the type and duration of surgery were significantly associated with POP ( $p$ -values were 0.002 and 0.006, respectively).

**Conclusions:** PONV and POP are common complications in our surgical patients. Factors associated with PONV include a prior history of PONV and POP. Patients at risk should be identified, the proper formulation of PONV protocols should be considered, and appropriate management plans should be implemented to improve patients' outcomes.

**Keywords:** Postoperative nausea and vomiting, Postoperative pain, Surgical patients

## Background

Efficacy of healthcare administration, raising the patient's level of satisfaction, and obtaining optimal outcomes are the major reflection tools for evaluating any health system's overall quality. Multiple postsurgical complications with different severities have been recorded, including

\*Correspondence: saedyoud@yahoo.com; saedyoud@najah.edu

<sup>3</sup> Department of Clinical and Community Pharmacy, Department of Pharmacy, College of Medicine and Health Sciences, An-Najah National University, 44839 Nablus, Palestine

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

postoperative nausea and vomiting (PONV) and perioperative pain [1, 2]. PONV is defined as the experience of nausea, vomiting, or both during the first 24 h following surgery. It is one of the most reported causes of distress and dissatisfaction among surgical patients in the early postoperative hours. It is frequently cited as one of the most unpleasant complications that patients would prefer to avoid, and successful prevention of this complication greatly improves their satisfaction and hastens their resumption of daily activities [3, 4].

PONV affects the decision of when to discharge patients from the recovery room and is associated with a delayed hospital stay, thus increasing ward burden and overall healthcare costs. Although typically self-limiting and rarely fatal, PONV increases the patients' risk of developing dehydration, serum electrolyte disturbances, aspiration and subsequent pneumonia, and, if severe enough, oesophageal rupture [5].

Recent studies showed that the incidence of PONV continues to be unacceptably high, ranging from 9 to 56% [6–8]. Furthermore, in high-risk patients undergoing surgery without receiving any prophylaxis, the incidence of PONV was reported to be as high as 80% [9, 10]. However, there is still a shortage of studies that discuss the pathophysiology and potential factors influencing the development of PONV, which might contribute to its high incidence rate [5, 11]. Many factors are proposed as risk factors for PONV. These can be categorized into patient-related and surgery-related factors. The Apfel score assesses four factors: sex, smoking status, postoperative opioid use, and prior motion sickness or PONV. Assessment of the patients' risk of experiencing PONV helps in predicting this problematic postoperative consequence [6].

Another highly reported complication in the postoperative period is pain [12]. Several studies have addressed this complication and helped improve our understanding of its underlying mechanisms. However, postoperative pain (POP) continues to be a prevalent yet unresolved concern [13, 14]. Recognizing factors associated with POP provides the basis for effective pain management and improved outcomes [15]. POP is also associated with other public health and economic concerns [13]. Therefore, effective pain management is also the key to achieving better outcomes on the societal level [16].

In Palestine, there is data on the incidence of PONV, the factors associated with it, and the association between PONV and POP. Evaluating pain and PONV is noteworthy because the key reasons for the failure of discharge after day-case surgery are pain and PONV. Therefore, this study aimed to determine whether PONV and POP were associated with demographic or surgical characteristics among patients recovering from surgery and

to evaluate the relationship between PONV and POP. The results of this study provide valuable data that can help in improving surgical outcomes for patients and the healthcare system at large.

## Methods

### Study design and setting

This was a prospective, multicentre, observational study conducted to estimate the incidence of PONV in surgical patients and their associations with different pain indices. The study was conducted at An-Najah National University Hospital, a large tertiary hospital and an important referral centre for patients from both the West Bank and Gaza, and Rafidia Governmental Hospital, a major surgical hospital in the northern West Bank. Both hospitals are in Nablus city in the West Bank, Palestine. Data was collected between October 2019 and February 2020. This study adhered to the STROBE guideline.

### Study area and population of the study

Our target population was surgical patients in Palestine. Participants were enrolled in this study based on the specified criteria for inclusion and exclusion.

### Sample size and sampling technique

A total of 211 patients undergoing elective surgery under general anaesthesia at An-Najah National University Hospital and Rafidia Surgical Governmental Hospital were selected using a convenient sampling procedure.

### Inclusion and exclusion criteria

Patients aged above 18 years old of both sexes who were scheduled for elective surgery under general anaesthesia and agreed to participate in our study were eligible to participate. Our exclusion criteria included admission to the intensive care unit (ICU), requiring the insertion of a nasogastric tube after surgery, having cognitive impairment, or receiving steroids or antiemetic medications in the last 24 h before surgery.

### Data collection instruments

We conducted data collection form-based interviews with the participants during the first 24 h following their surgery. The data collection form consisted of four sections [17–19] (Additional file 1: Data collection form). The first section contained questions inquiring about the participant's demographic (age and sex) and clinical (BMI, smoking, history of motion sickness and history of PONV) characteristics. The second section inquired about their current operation, including the type of surgery, duration of surgery, postoperative opioid use and POP.

The third section assessed postoperative nausea (defined as pronounced stomach discomfort with the sensation of an urge to vomit) using a visual analogue scale (VAS) that was approved for assessing the intensity of nausea in the postoperative period [17]. Participants rated the overall intensity of nausea they felt during the first 24 h of the postoperative period on a 100-mm VAS, with 0 meaning no nausea at all and 100 meaning severe nausea. In this section, we also collected data on postoperative vomiting by asking about the number of vomiting episodes experienced in the 24 h following surgery. PONV, defined as suffering from postoperative nausea, vomiting, or both, was also assessed in this section.

The last section contained items that assessed POP severity during the first 24 h following surgery using a numerical rating scale [18]. Participants reported their response as a number from 0 to 10, with 0 meaning no pain at all, 1–3 meaning mild pain, 4–6 meaning moderate pain and 7–10 meaning severe pain. We recorded the severity of the average overall pain, as well as the severity of pain on movement (defined as pain experienced with any active movement, including getting out of bed, sitting, turning to the side, defecating, taking deep breaths and coughing forcefully) and pain at rest (defined as pain occurring while in bed with no active movements).

Data on anaesthetic drugs and analgesic administration was collected from the intraoperative anaesthetic charts and the inpatients electronic recording system after obtaining consent to use this data from both the hospital and the participants.

The data collection form was pilot-tested on 20 patients at An-Najah National University Hospital.

### Ethical approval

The study was reviewed and approved by the *Institutional Review Board (IRB) of An-Najah National University*. Participants had the right to choose whether they would like to be included in the study or not. Verbal consent was obtained, and this was followed by providing the participant with written informed consent to read and sign voluntarily.

### Statistical analysis

All statistical analysis was conducted using version 21.0 of Statistical Package for Social Sciences (SPSS) (IBM Corp., Armonk, NY, USA). Variables were presented as frequencies and percentages and/or medians and interquartile ranges, as appropriate. Pearson's chi-square test was used for correlational analysis between variables, and p-values were reported. For PONV assessment, we divided our participants into two groups: those who had PONV and those who did not have PONV. All univariate variables significant at  $p < 0.05$

were entered into a multiple binary logistic regression model to determine factors that were significantly associated with the occurrence of PONV. By estimating the odds ratio (OR) at a 95% confidence interval (CI), the association between the occurrence of PONV and each independent variable was measured. For pain assessment, patients were categorized into two groups: those who had POP and those who did not have POP. A p-value less than 0.05 was considered statistically significant.

## Results

### Demographic and clinical characteristics and their association with PONV

Out of the 221 subjects approached, 211 were included in this study, accounting for a response rate of 95.5%. The demographic and clinical characteristics of the participants are presented in Table 1. The median age of the participants was 33 years, with an interquartile range (IQR) of 24–48. The male-to-female ratio was roughly balanced (55.5% and 44.5%, respectively). Almost half (46.4%) of the participants were overweight (BMI in the 25–29.9 range) and the majority had no history of motion sickness (87.7%) or PONV (82.5%). Figure 1 shows a flow diagram of the study patients. Using the Apfel simplified score, participants can be categorized depending on the number of risk factors they have, into 0, 1, 2, 3 or 4 risk factors, as shown in Table 2. Most of the participants had only 1 or 2 risk factors (30.3% and 31.3%, respectively). Of the 211 participants, 43.1% reported experiencing nausea, whereas only 17.5% reported having vomited. The overall incidence of PONV in our sample was 45.5% (Table 2).

Reported PONV was significantly correlated with a history of PONV among the participants ( $p = 0.009$ ). No other significant association was observed between reported PONV and demographic or clinical characteristics (Table 1).

### Operation-related factors and their association with PONV


The most frequent types of surgery that participants underwent were general surgery/laparotomy (30.8%), laparoscopic surgery (21.3%) and orthopaedic surgery (12.3%). Table 3 presents the percentage of participants based on the type of surgery they underwent and shows the association of PONV with operation-related factors among the participants. The type of surgery showed a significant association with PONV ( $p = 0.010$ ). Additionally, we found a significant association between the occurrence of PONV and experiencing POP ( $p = 0.002$ ).

**Table 1** Demographic and clinical characteristics and their relationship with PONV

| Characteristic             | Total; N = 211 (%) | PONV; N = 96 (%) | No PONV; N = 115 (%) | P-value*     |
|----------------------------|--------------------|------------------|----------------------|--------------|
| Age (year)                 |                    |                  |                      |              |
| 18–39                      | 135 (64)           | 59 (61.5)        | 76 (66.1)            | 0.658        |
| 40–64                      | 66 (31.3)          | 33 (34.4)        | 33 (28.7)            |              |
| 65–79                      | 10 (4.7)           | 4 (4.2)          | 6 (5.2)              |              |
| Sex                        |                    |                  |                      |              |
| Female                     | 94 (44.5)          | 48 (50.0)        | 46 (40)              | 0.146        |
| Male                       | 117 (55.5)         | 48 (50.0)        | 69 (60)              |              |
| BMI                        |                    |                  |                      |              |
| Underweight                | 4 (1.9)            | 3 (3.1)          | 1 (0.9)              | 0.208        |
| Healthy weight             | 57 (27.0)          | 21 (21.9)        | 36 (31.3)            |              |
| Overweight                 | 98 (46.4)          | 50 (52.1)        | 48 (41.7)            |              |
| Obese                      | 52 (24.6)          | 22 (22.9)        | 30 (26.1)            |              |
| Smoking                    | 93 (44.1)          | 40 (41.7)        | 53 (46.1)            | 0.520        |
| History of motion sickness | 26 (12.3)          | 10 (10.4)        | 16 (13.9)            | 0.442        |
| History of PONV            | 37 (17.5)          | 24 (25.0)        | 13 (11.3)            | <b>0.009</b> |

PONV postoperative nausea and vomiting, BMI body mass index

\*Significant p-values are in bold



**Fig. 1** Flow diagram of the study patients. PONV postoperative nausea and vomiting, POP postoperative pain

**Occurrence and severity of POP and its association with PONV**

Most participants (74.4%) reported experiencing some degree of POP (Table 3), with 14.2% experiencing severe pain at rest and 29.4% experiencing severe pain with movement. Data on the occurrence and severity of POP, in general and per set, is presented in Table 4. Of all

participants with POP, 46.9% underwent short operations (surgery duration of <60 min). There were also significant associations between PONV and pain severity in general, at rest, and with movement (p-values were <0.001, <0.001, and 0.010, respectively), as shown in Table 4. Both the type and duration of surgery were significantly correlated with the occurrence of POP

**Table 2** Apfel score risk factors

| Apfel score   | Frequency | PONV; N=96 | No PONV; N=115 | P value |
|---------------|-----------|------------|----------------|---------|
| Apfel score=0 | 46 (21.8) | 17 (17.7)  | 29 (25.2)      | 0.240   |
| Apfel score=1 | 64 (30.3) | 28 (29.2)  | 36 (31.3)      |         |
| Apfel score=2 | 66 (31.3) | 31 (32.3)  | 35 (30.4)      |         |
| Apfel score=3 | 29 (13.7) | 15 (15.6)  | 14 (12.2)      |         |
| Apfel score=4 | 6 (2.8)   | 5 (5.2)    | 1 (0.9)        |         |
| Overall       | 211 (100) | 96 (100)   | 115 (100)      |         |

(*p*-values were 0.002 and 0.006, respectively). Table 5 shows the association of POP with operation-related factors among the participants.

### Use of drugs and their association with both PONV and pain

Despite the diversity of operations that our participants underwent, almost all participants received the same anaesthetic drugs. Propofol, fentanyl, midazolam and atracurium were a standardized regimen. The rates of PONV in association with these drugs were 100%, 99%, 38.5% and 53.1%, respectively. Table 6 shows the percentage of usage of these drugs, as well as the frequency of experiencing PONV and POP. Most (69.4%) of the patients who reported experiencing pain in the postoperative period were given opioids. Among those, 28.1% experienced PONV, whereas 71.9% of the 161 patients (76.3% of all participants) who were managed with non-opioid analgesics (mainly acetaminophen) had PONV.

### Multiple logistic regression analysis

Variables with a *p*-value < 0.05, including the history of PONV, POP and type of surgery, were entered in a multiple binary logistic regression model. Some of the above associations did not exist after controlling other variables. In multiple binary logistic regression analysis using PONV as a dependent variable, only patients with a history of PONV (OR = 2.28; 95% CI = 1.03–5.01; *p* = 0.041) and POP (OR = 2.41; 95% CI = 1.17–4.97; *p* = 0.018) were significantly associated with the occurrence of PONV (Table 7). All included variables in the multiple binary logistic regression had a significant *p*-value in the univariate analysis between PONV and non-PONV groups. The model was significant, with a Chi-square of 28.67, DF = 8; *p* < 0.001.

### Discussion

In the current study, we assessed the incidence of PONV and POP, their association, and other factors that could predict their occurrence in surgical patients. This was the first study of its kind in Palestine. We found that PONV and POP are common complications in the first 24 h following surgery. We also reported a significant association between these two complications, as well as their association with other factors. PONV was associated with previous PONV and surgery type, whereas POP was associated with surgery duration.

PONV and POP significantly affect morbidity and patient satisfaction in the immediate postoperative period. The overall incidence of PONV, as described in our study, is comparable to the findings of a 2012 systematic review that included 22 studies [6].

**Table 3** Operation-related factors and their association with PONV

| Characteristic        | Total; N = 211 (%) | PONV; N = 96 (%) | No PONV; N = 115 (%) | P-value*     |
|-----------------------|--------------------|------------------|----------------------|--------------|
| Type of surgery       |                    |                  |                      |              |
| General               | 65 (30.8)          | 30 (31.3)        | 35 (30.4)            | <b>0.010</b> |
| Laparoscopic          | 45 (21.3)          | 26 (27.1)        | 19 (16.5)            |              |
| Orthopedic            | 26 (12.3)          | 15 (15.6)        | 11 (9.6)             |              |
| Otorhinolaryngology   | 35 (16.6)          | 11 (11.5)        | 24 (20.9)            |              |
| Urosurgery            | 21 (10.0)          | 5 (5.2)          | 16 (13.9)            |              |
| Neurosurgery          | 11 (5.2)           | 8 (8.3)          | 3 (2.6)              |              |
| Vascular surgery      | 4 (1.9)            | 1 (1.0)          | 3 (2.6)              |              |
| Gynecological surgery | 4 (1.9)            | 0 (0.0)          | 4 (3.5)              |              |
| Duration of surgery   |                    |                  |                      |              |
| < 60 min              | 114 (68.2)         | 63 (65.6)        | 81 (70.4)            | 0.455        |
| ≥ 60 min              | 67 (31.8)          | 33 (34.4)        | 34 (29.6)            |              |
| Post-operative opioid | 50 (23.7)          | 27 (28.1)        | 23 (20.0)            | 0.167        |
| Post-operative pain   | 157 (74.4)         | 81 (84.4)        | 76 (66.1)            | <b>0.002</b> |

PONV postoperative nausea and vomiting, BMI body mass index

\*Significant *p*-values are in bold



**Table 4** The occurrence and severity of postoperative pain and their association with PONV

| Characteristic            | Frequency (%) | PONV; N = 96 (%) | No PONV; N = 115 (%) | P-value*          |
|---------------------------|---------------|------------------|----------------------|-------------------|
| <b>Pain severity</b>      |               |                  |                      |                   |
| None                      | 54 (25.6)     | 15 (27.8)        | 39 (72.2)            | <b>&lt; 0.001</b> |
| Mild                      | 56 (26.5)     | 22 (39.3)        | 34 (60.7)            |                   |
| Moderate                  | 64 (30.3)     | 31 (48.4)        | 33 (51.6)            |                   |
| Severe                    | 37 (17.5)     | 28 (75.7)        | 9 (24.3)             |                   |
| <b>Pain at rest</b>       |               |                  |                      |                   |
| None                      | 54 (25.6)     | 15 (27.8)        | 39 (72.7)            | <b>&lt; 0.001</b> |
| Mild                      | 72 (34.1)     | 34 (47.2)        | 38 (52.8)            |                   |
| Moderate                  | 55 (26.1)     | 23 (41.8)        | 32 (58.2)            |                   |
| Severe                    | 30 (14.2)     | 24 (80.0)        | 6 (20.0)             |                   |
| <b>Pain with movement</b> |               |                  |                      |                   |
| None                      | 54 (25.6)     | 15 (27.8)        | 39 (72.2)            | <b>0.010</b>      |
| Mild                      | 56 (26.5)     | 25 (44.6)        | 31 (55.4)            |                   |
| Moderate                  | 39 (18.5)     | 20 (51.3)        | 19 (48.7)            |                   |
| Severe                    | 62 (29.4)     | 36 (58.1)        | 26 (41.9)            |                   |

PONV postoperative nausea and vomiting

\*Significant p-values are in bold

Factors that showed a statistically significant association with PONV in the 2012 review included previous PONV, which was an association that we also found to be significant in our sample. It is postulated that a history of PONV points to the presence of an underlying susceptibility to PONV [6]. This finding was also in agreement with the findings of other studies that were conducted in Ethiopia [20], Korea [21] and Uganda [22].

A possible explanation for this finding was suggested by a 2011 study that used pooled DNA samples to detect genetic markers that would potentially influence the possibility of experiencing PONV [23]. In that study, subjects with previous PONV were more likely to have first-degree relatives with a history of PONV [23].

The presence of motion sickness history, on the other hand, did not have a significant effect on experiencing PONV in our study, similar to the findings of a study that examined 174 patients undergoing minor orthopaedic surgery [24]. Another more recent study reported similar findings on the relationship between a history of motion sickness and PONV [22].

**Table 5** Operation-related factors and their association with pain occurrence

| Operation-related factors  | With post-operative pain; N = 157 (%) | Without post-operative pain; N = 54 (%) | P value*     |
|----------------------------|---------------------------------------|---|--------------|
| <b>Type of surgery</b>     |                                       |   |              |
| General surgery/Laparotomy | 51 (32.5)                             | 14(25.9)                                | <b>0.002</b> |
| Laparoscopic               | 38 (24.2)                             | 7 (13.0)                                |              |
| Orthopedic                 | 22 (14.0)                             | 4 (7.4)                                 |              |
| Otorhinolaryngology        | 24 (15.3)                             | 11 (20.4)                               |              |
| Urosurgery                 | 9 (5.7)                               | 12 (22.2)                               |              |
| Neurosurgery               | 8 (5.1)                               | 3 (5.6)                                 |              |
| Vascular                   | 1 (0.5)                               | 3 (5.6)                                 |              |
| Gynecological              | 4 (1.9)                               | 0 (0.0)                                 |              |
| <b>Duration of surgery</b> |                                       |   |              |
| < 60 min                   | 99 (63.1)                             | 45 (83.3)                               | <b>0.006</b> |
| ≥ 60 min                   | 58 (36.9)                             | 9 (16.7)                                |              |

\*Significant p-values are in bold

**Table 6** Association between PONV, pain and the drugs used

| Medications | PONV; N = 96 (%) | No PONV; N = 115 (%) | P value* | Pain; N = 157 | No pain; N = 54 | P value*     |
|-------------|------------------|----------------------|----------|---------------|-----------------|--------------|
| Propofol    | 96 (100)         | 113 (98.3)           | 0.296    | 155 (98.7)    | 54 (100)        | 0.553        |
| Fentanyl    | 95 (99)          | 113 (98.3)           | 0.568    | 156 (99.4)    | 52 (96.3)       | 0.162        |
| Midazolam   | 37 (38.5)        | 42 (36.5)            | 0.436    | 60 (38.2)     | 19 (35.2)       | 0.410        |
| Atracurium  | 51 (53.1)        | 58 (50.4)            | 0.401    | 83 (52.9)     | 26 (48.1)       | 0.330        |
| Opioid      | 27 (28.1)        | 23 (20)              | 0.111    | 109 (69.4)    | 52 (96.3)       | <b>0.000</b> |

\*Significant p-values are in bold

**Table 7** Independent factors associated with PONV using multiple logistic regression analysis (enter method)

| Variable            | B      | S.E  | Wald | P value* | Odds ratio with 95% CI |
|---------------------|--------|------|------|----------|------------------------|
| History of PONV     |        |      |      |          |                        |
| No                  |        |      |      |          | Ref                    |
| Yes                 | 0.82   | 0.40 | 4.18 | 0.041    | 2.28 (1.03–5.01)       |
| Post-operative pain |        |      |      |          |                        |
| No                  |        |      |      |          | Ref                    |
| Yes                 | 0.88   | 0.37 | 5.64 | 0.018    | 2.41 (1.17–4.97)       |
| Type of surgery     |        |      |      |          |                        |
| General             |        |      |      |          | Ref                    |
| Laparoscopic        | 0.52   | 0.40 | 1.64 | 0.200    | 1.68 (0.76–3.69)       |
| Orthopedic          | 0.40   | 0.48 | 0.77 | 0.401    | 1.50 (0.58–3.85)       |
| Otorhinolaryngology | − 0.48 | 0.45 | 1.11 | 0.292    | 0.62 (0.26–1.51)       |
| Urosurgery          | − 0.59 | 0.59 | 0.97 | 0.324    | 0.56 (0.17–1.78)       |
| Neurosurgery        | 1.16   | 0.75 | 2.40 | 0.122    | 3.17 (0.74–13.67)      |
| Vascular surgery    | − 1.52 | 1.11 | 1.86 | 0.172    | 0.22 (0.26–1.94)       |

PONV postoperative nausea and vomiting, CI confidence interval, B the coefficient of the predictor variables

\*Significant p-values are in bold

Our results also showed that the type of operative procedure and the incidence of PONV were both significantly associated with experiencing PONV. Similar findings were also reported by a study conducted on 17,638 surgical patients to detect predictors of PONV [25] and by another study that was designed to examine the influence of the type of surgery on the occurrence of PONV [26].

Neither age nor sex was a significant predictor of the occurrence of PONV in this study, similar to the results of a study conducted in Ghana [27]. Another similar prospective, interview-based survey conducted on 1,107 patients between 4 and 86 years old did not find any effect of age on the occurrence of nausea [28]. Furthermore, one study reported an equal incidence of PONV in both males and females undergoing extra-abdominal operations [29]. Although the female sex was found to be a strong risk factor for PONV in some studies, this effect might or might not be mediated by other factors and requires more investigation. For example, it has been proposed that serum progesterone levels, the day of the menstrual cycle, and menopause might have a confounding effect [27]. A retrospective review that examined the influence of the menstrual cycle on PONV found that the incidence was at its highest on day five of the cycle and that there was no nausea and vomiting on days 18–20 [30]. Ethnicity may also have a major role in explaining these results. A prospective study that was conducted to

compare the incidence of PONV in different ethnicities concluded that ethnicity could be an independent risk factor for developing PONV [31].

Although non-smoking status and postoperative opioid use were among the predictors of PONV in Apfel's predictive model [19], no such associations were significant in the current study, similar to other studies from Ethiopia [20] and Uganda [22]. Furthermore, a study that assessed PONV in patients undergoing arthroscopic surgery concluded that short-acting opioids did not have a significant effect on patients' risk of developing PONV [32]. These findings suggest that the effect of smoking and opioid use on PONV might vary, depending on other patient- and surgery-related factors, and should be investigated further. For example, previous studies have found that the type of surgery might affect the incidence of vomiting upon changing the route of administration in patients receiving morphine, and that vomiting might occur more frequently in patients undergoing abdominal operations compared to hip replacement procedures and in patients receiving epidural opioids compared to intravenous patient-controlled analgesia [33, 34]. Another study reported that intravenous injection of morphine or ketobemidone relieved 80% of nausea episodes [34]. It was also reported that opioid use could rarely lead to postoperative nausea in immobile patients [35].

A high number of patients in this study reported moderate to severe pain during the first 24 h following surgery, similar to findings reported by other studies [8, 11, 36]. Additionally, POP was significantly associated with the development of PONV, which is also similar to reports in the literature [34, 35]. For example, a prospective study that aimed to examine predictors of PONV reported a higher frequency of PONV among patients experiencing excessive POP [25]. A significant association between reported pain and PONV was also seen in several other studies [26, 37, 38]. A study conducted in Uganda found that surgical patients who reported feeling pain were twice as likely to experience PONV than patients who did not [22]. It was also noted that nausea was frequently accompanied by pain in the first few hours after surgery, and that relieving the pain led to a decrease in nausea [25, 35]. Similarly, a study conducted in Norway found that managing pain reduced nausea symptoms [35]. These findings are in agreement with findings from previous studies that concluded that the use of intravenous acetaminophen postoperatively might reduce PONV incidence and the need to give antiemetic medications [39, 40].

Several methods have been used in the recovery room to minimize POP. Opioids continue to be a popular means, despite their widely recognized adverse effects.

Recently, non-opioid analgesics and adjuncts, such as dexamethasone, have been used as opioid-sparing alternatives [41]. Recent guidelines recommend using second-generation 5-hydroxytryptamine 3 (5-HT<sub>3</sub>) receptor antagonists, dopamine antagonists and neurokinin 1 (NK1) receptor antagonists as prophylaxis and treatment options for nausea and/or vomiting in the postoperative setting [42].

In summary, the current study found a high PONV incidence among surgical patients and identified factors associated with PONV, including previous PONV, type of surgery and POP. Additionally, we found a high incidence of POP, necessitating the implementation of appropriate pain management protocols during the postoperative period.

### Strengths and limitations

The current study had some limitations. Participants were interviewed within 24 h following their surgery, which may have limited our findings to symptoms occurring inside this time window. Moreover, our study did not include PONV and POP prophylaxis and management protocols followed by the hospitals studied. Furthermore, the high variability of the population selected (e.g., pain, opioid consumption, type of surgery, Apfel risk factors, etc.) might interfere with the statistical analysis results. The final limitation is the lack of information regarding pain management methods, including whether the subjects received regional, local or parenteral analgesia. Thus, we are unable to assess if the methods of analgesia affect the occurrence of POP. On the other hand, this study had many strengths. This was the first study to assess PONV, POP, and their association in Palestine. We also sampled our participants from two of the largest surgical hospitals in the country, which may have improved the generalizability of our findings.

### Conclusions

In conclusion, our findings suggest that PONV and POP are common complications in our surgical patients. Risk factors of PONV include prior history of PONV and POP. Patients at risk should be identified, the proper formulation of PONV protocols should be considered, and appropriate POP management should be highlighted and reinforced to improve patients' outcomes during the postoperative period.

### Abbreviations

BMI: Body mass index; CI: Confidence interval; ICU: Intensive care unit; IQR: Interquartile range; IRB: Institutional review board; OR: Odds ratio; PONV: Postoperative nausea and vomiting; POP: Postoperative pain; VAS: Visual analogue scale.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12893-021-01172-9>.

**Additional file 1:** Data collection form. This is the final version of the English version that was used to obtain data to evaluate the relationship between postoperative nausea and vomiting and postoperative pain, and to identify some factors associated with these symptoms.

### Acknowledgements

Not applicable.

### Authors' contributions

RME and ASN performed data collection and analysis, reviewed the literature and contributed to drafting the manuscript. AS performed data analysis, reviewed the literature, contributed to data interpretation and assisted in drafting the manuscript. WS contributed to the conception and design of the study and interpreting the data. SZ conceptualized and designed the study, coordinated, supervised, and was responsible for the data integrity and analysis accuracy, critically reviewed the interpretation of the results and assisted in the final write-up. All authors read and approved the final manuscript.

### Funding

None.

### Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### Declarations

#### Ethics approval and consent to participate

The study was reviewed and approved by the *Institutional Review Board (IRB) of An-Najah National University*. Participants had the right to choose whether they would like to be included in the study or not. All participants were provided with written informed consent to read and sign voluntarily. Each participant gave written informed consent, which was approved by the IRB, to be included in the study, with the anonymous publication of data.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup> Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, 44839 Nablus, Palestine. <sup>2</sup> Department of Anesthesia and Intensive Care, An-Najah National University Hospital, 44839 Nablus, Palestine. <sup>3</sup> Department of Clinical and Community Pharmacy, Department of Pharmacy, College of Medicine and Health Sciences, An-Najah National University, 44839 Nablus, Palestine. <sup>4</sup> Poison Control and Drug Information Center (PCDIC), College of Medicine and Health Sciences, An-Najah National University, 44839 Nablus, Palestine. <sup>5</sup> Clinical Research Center, An-Najah National University Hospital, 44839 Nablus, Palestine.

Received: 7 July 2020 Accepted: 22 March 2021

Published online: 01 April 2021

### References

1. Aubrun F, Ecoffey C, Benhamou D, Jouffroy L, Diemunsch P, Skaare K, Bosson JL, Albaladejo P. Perioperative pain and post-operative nausea and vomiting (PONV) management after day-case surgery: the SFAR-OPERA national study. *Anaesth Crit Care Pain Med*. 2019;38(3):223–9.
2. Odom-Forren J, Rayens MK, Gokun Y, Jalota L, Radke O, Hooper V, Wiggins AT, Apfel CC. The relationship of pain and nausea in



- postoperative patients for 1 week after ambulatory surgery. *Clin J Pain*. 2015;31(10):845–51.
3. Fero KE, Jalota L, Hornuss C, Apfel CC. Pharmacologic management of postoperative nausea and vomiting. *Expert Opin Pharmacother*. 2011;12(15):2283–96.
  4. Pierre S, Whelan R. Nausea and vomiting after surgery. *Cont Educ Anaesthesia Crit Care Pain*. 2013;13(1):28–32.
  5. Shaikh SI, Nagarekha D, Hegade G, Marutheesh M. Postoperative nausea and vomiting: a simple yet complex problem. *Anesth Essays Res*. 2016;10(3):388–96.
  6. Apfel CC, Heidrich FM, Jukar-Rao S, Jalota L, Hornuss C, Whelan RP, Zhang K, Cakmakaya OS. Evidence-based analysis of risk factors for postoperative nausea and vomiting. *Br J Anaesth*. 2012;109(5):742–53.
  7. Amirshahi M, Behnamfar N, Badakhsh M, Rafiemanesh H, Keikhaie KR, Sheyback M, Sari M. Prevalence of postoperative nausea and vomiting: a systematic review and meta-analysis. *Saudi J Anaesth*. 2020;14(1):48–56.
  8. Dolin SJ, Cashman JN, Bland JM. Effectiveness of acute postoperative pain management: I. Evidence from published data. *Br J Anaesth*. 2002;89(3):409–23.
  9. Eberhart LH, Frank S, Lange H, Morin AM, Scherag A, Wulf H, Kranke P. Systematic review on the recurrence of postoperative nausea and vomiting after a first episode in the recovery room—implications for the treatment of PONV and related clinical trials. *BMC Anesthesiol*. 2006;6:14.
  10. Kovac AL. Update on the management of postoperative nausea and vomiting. *Drugs*. 2013;73(14):1525–47.
  11. Lee YZ, Lee RQ, Thinn KK, Poon KH, Liu EH. How patients fare after anaesthesia for elective surgery: a survey of postoperative nausea and vomiting, pain and confusion. *Singapore Med J*. 2015;56(1):40–6.
  12. Gupta A, Kaur K, Sharma S, Goyal S, Arora S, Murthy RS. Clinical aspects of acute post-operative pain management & its assessment. *J Adv Pharm Technol Res*. 2010;1(2):97–108.
  13. Gan TJ. Poorly controlled postoperative pain: prevalence, consequences, and prevention. *J Pain Res*. 2017;10:2287–98.
  14. Joshi GP, Ogunnaik BO. Consequences of inadequate postoperative pain relief and chronic persistent postoperative pain. *Anesthesiol Clin North Am*. 2005;23(1):21–36.
  15. Mei W, Seeling M, Franck M, Radtke F, Brantner B, Wernecke KD, Spies C. Independent risk factors for postoperative pain in need of intervention early after awakening from general anaesthesia. *Eur J Pain*. 2010;14(2):149.
  16. Sinatra R. Causes and consequences of inadequate management of acute pain. *Pain Med*. 2010;11(12):1859–71.
  17. Boogaerts JG, Vanacker E, Seidel L, Albert A, Bardiau FM. Assessment of postoperative nausea using a visual analogue scale. *Acta Anaesthesiol Scand*. 2000;44(4):470–4.
  18. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EK, Kvarstein G, Stubhaug A. Assessment of pain. *Br J Anaesth*. 2008;101(1):17–24.
  19. Weillbach C, Rahe-meyer N, Raymondos K, Weissig A, Scheinichen D, Piepenbrock S. Postoperative nausea and vomiting (PONV): usefulness of the Apfel-score for identification of high risk patients for PONV. *Acta Anaesthesiol Belg*. 2006;57(4):361–3.
  20. Ahmed SA, Lema GF. Incidence and factors associated with postoperative nausea and vomiting among elective adult surgical patients at University of Gondar comprehensive specialized hospital, Northwest Ethiopia, 2019: a cross-sectional study. *Int J Surg Open*. 2020;22:57–61.
  21. Yi MS, Kang H, Kim MK, Choi GJ, Park YH, Baek CW, Jung YH, Woo YC. Relationship between the incidence and risk factors of postoperative nausea and vomiting in patients with intravenous patient-controlled analgesia. *Asian J Surg*. 2018;41(4):301–6.
  22. Ssebuufu R, Kakande I, Okello M. Post-operative nausea and vomiting at Mulago Hospital. *East Cent African J Surg*. 2009;14(2):50–7.
  23. Janicki PK, Vealey R, Liu J, Escajeda J, Postula M, Welker K. Genome-wide Association study using pooled DNA to identify candidate markers mediating susceptibility to postoperative nausea and vomiting. *Anesthesiology*. 2011;115(1):54–64.
  24. Tramer MR. A rational approach to the control of postoperative nausea and vomiting: evidence from systematic reviews. Part I. Efficacy and harm of antiemetic interventions, and methodological issues. *Acta Anaesthesiol Scand*. 2001;45(1):4–13.
  25. Sinclair DR, Chung F, Mezei G. Can postoperative nausea and vomiting be predicted? *Anesthesiology*. 1999;91(1):109–18.
  26. Doubravska L, Dostalova K, Fritscherova S, Zapletalova J, Adamus M. Incidence of postoperative nausea and vomiting in patients at a university hospital. Where are we today? *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2010;154(1):69–76.
  27. Bellville JW, Bross ID, Howland WS. Postoperative nausea and vomiting. IV. Factors related to postoperative nausea and vomiting. *Anesthesiology*. 1960;21:186–93.
  28. Koivuranta M, Laara E, Snare L, Alahuhta S. A survey of postoperative nausea and vomiting. *Anaesthesia*. 1997;52(5):443–9.
  29. Dyrberg V. Haloperidol (Serenase) in the prevention of postoperative nausea and vomiting. *Acta Anaesthesiol Scand*. 1962;6:37–47.
  30. Beattie WS, Lindblad T, Buckley DN, Forrest JB. The incidence of postoperative nausea and vomiting in women undergoing laparoscopy is influenced by the day of menstrual cycle. *Can J Anaesth*. 1991;38(3):298–302.
  31. Alli A, Omar S, Tsang S, Naik BI. The Effect of ethnicity on the incidence of postoperative nausea and vomiting in moderate to high risk patients undergoing general anaesthesia in South Africa: a controlled observational study. *Middle East J Anaesthesiol*. 2017;24(2):119–29.
  32. Dershwitz M, Michalowski P, Chang Y, Rosow CE, Conlay LA. Postoperative nausea and vomiting after total intravenous anesthesia with propofol and remifentanyl or alfentanil: how important is the opioid? *J Clin Anesth*. 2002;14(4):275–8.
  33. Weller R, Rosenblum M, Conard P, Gross JB. Comparison of epidural and patient-controlled intravenous morphine following joint replacement surgery. *Can J Anaesth*. 1991;38(5):582–6.
  34. Egbert AM, Parks LH, Short LM, Burnett ML. Randomized trial of postoperative patient-controlled analgesia vs intramuscular narcotics in frail elderly men. *Arch Intern Med*. 1990;150(9):1897–903.
  35. Andersen R, Krohg K. Pain as a major cause of postoperative nausea. *Can Anaesth Soc J*. 1976;23(4):366–9.
  36. Sommer M, de Rijke JM, van Kleef M, Kessels AG, Peters ML, Geurts JW, Gramke HF, Marcus MA. The prevalence of postoperative pain in a sample of 1490 surgical inpatients. *Eur J Anaesthesiol*. 2008;25(4):267–74.
  37. Quinn AC, Brown JH, Wallace PG, Asbury AJ. Studies in postoperative sequelae. Nausea and vomiting—still a problem. *Anaesthesia*. 1994;49(1):62–5.
  38. Urits I, Orhurhu V, Jones MR, Adamian L, Borchart M, Galasso A, Viswanath O. Postoperative nausea and vomiting in paediatric anaesthesia. *Turk J Anaesthesiol Reanim*. 2020;48(2):88–95.
  39. Taniguchi H, Nagahuchi M, Kamada T, Suga K, Sato T, Ueda T, Yamamoto T. Effect of scheduled intravenous acetaminophen on postoperative nausea and vomiting in patients undergoing laparoscopic gynecologic surgery. *J Anesth* 2020.
  40. Watcha MF, White PF. Postoperative nausea and vomiting. Its etiology, treatment, and prevention. *Anesthesiology*. 1992;77(1):162–84.
  41. Luo J, Min S. Postoperative pain management in the postanesthesia care unit: an update. *J Pain Res*. 2017;10:2687–98.
  42. Gan TJ, Belani KG, Bergese S, Chung F, Diemunsch P, Habib AS, Jin Z, Kovac AL, Meyer TA, Urman RD, et al. Fourth consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg*. 2020;131(2):411–48.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.